

ELENCO 10 PUBBLICAZIONI PIÙ SIGNIFICATIVE - F.M. RIJLI

Kitazawa T, Machlab D, Joshi O, Maiorano NA, Kohler H, Ducret S, Gezelius H, Soneson C, Papasaikas P, López-Bendito G, Stadler MB, **Rijli FM** (2021) A unique bipartite Polycomb signature regulates stimulus response transcription during development. *Nature Genetics*, 53(3):379-391.

Minoux M, Holwerda S, Vitobello A, Kitazawa T, Kohler H, Stadler M, **Rijli FM** (2017) Gene Bivalency at Polycomb Domains Regulates Cranial Neural Crest Positional Identity. *Science*, 355(6332), eaal2913. (full article)

Di Meglio T, Kratochwil CF, Vilain N, Loche A, Vitobello A, Yonehara K, Hrycaj SM, Roska B, Peters AH, Eichmann A, Wellik D, Ducret S, **Rijli FM** (2013) Ezh2 orchestrates topographic tangential migration and connectivity of precerebellar neurons. *Science*, 339(6116):204-7.

Vitobello A, Ferretti E, Lampe X, Vilain N, Ducret S, Ori M, Spetz JF, Selleri L, **Rijli FM** (2011) Hox and Pbx factors control retinoic acid synthesis during hindbrain segmentation. *Developmental Cell*, 20(4):469-82.

Erzurumlu, R., Murakami, Y., and **Rijli F.M.** (2010) Mapping the face in the somatosensory brainstem. *Nature Reviews Neuroscience*, 11(4):252-63.

Oury, F., Murakami, Y., Renaud, J-S, Pasqualetti, M., Charnay, P., Ren, S-Y., and **Rijli F.M.** (2006) Hoxa2 and rhombomere dependent development of the mouse facial somatosensory map. *Science*, 313(5792):1408-13. (full article)

Santagati, F., and **Rijli F.M.** (2003). Cranial neural crest and the building of the vertebrate head. *Nature Reviews Neuroscience*, 4, 806–818.

Pasqualetti M., Neun, R., Davenne M., and **Rijli F.M.** (2001) Retinoic acid rescues inner ear defects in Hoxa1 deficient newborn mice. *Nature Genetics*, 29(1):34-9. (full article)

Davenne, M., Maconochie, M.K., Neun, R., Pattyn, A., Chambon, P., Krumlauf, R., and **Rijli, F.M.** (1999) Hoxa2 and Hoxb2 control dorsoventral patterns of neuronal development in the rostral hindbrain. *Neuron*, 22, 677-691.

Rijli, F.M., Mark, M., Lakkaraju, S., Dierich, A., Dollé, P., and Chambon, P. (1993). A homeotic transformation is generated in the rostral branchial region of the head by disruption of Hoxa-2, which acts as a selector gene. *Cell*, 75, 1333-1349.

BIBLIOGRAFIA COMPLETA- F.M. RIJLI

Genescu I, Aníbal-Martínez M, Kouskoff V, Chenouard N, Mailhes-Hamon C, Cartonnet H, Lokmane L, **Rijli FM**, López-Bendito G, Gambino F, Garel S (2022) Dynamic interplay between thalamic activity and Cajal-Retzius cells regulates the wiring of cortical layer 1. *Cell Reports*, 12;39(2):110667.

Pumo GM, Kitazawa T, **Rijli FM** (2022) Epigenetic and Transcriptional Regulation of Spontaneous and Sensory Activity Dependent Programs During Neuronal Circuit Development. *Front Neural Circuits* 16:911023.

Machlab D, Burger L, Soneson C, **Rijli FM**, Schübeler D, Stadler MB (2022) monaLisa: an R/Bioconductor package for identifying regulatory motifs. *Bioinformatics*, 38(9):2624-5.

Kitazawa T, Minoux M, Ducret S, **Rijli FM** (2022) Different Ectopic Hoxa2 Expression Levels in Mouse Cranial Neural Crest Cells Result in Distinct Craniofacial Anomalies and Homeotic Phenotypes. *J. Dev. Biol.*, 10(1): 9.

Kitazawa T, Machlab D, Joshi O, Maiorano NA, Kohler H, Ducret S, Gezelius H, Soneson C, Papasaikas P, López-Bendito G, Stadler MB, **Rijli FM** (2021) A unique bipartite Polycomb signature regulates stimulus response transcription during development. *Nature Genetics*, Mar;53(3):379-391.

Maheshwari U, Kraus D, Vilain N, Holwerda SJB, Cankovic V, Maiorano NA, Kohler H, Satoh D, Sigrist M, Arber S, Kratochwil CF, Di Meglio T, Ducret S, **Rijli FM** (2020) Postmitotic Hoxa5 Expression Specifies Pontine Neuron Positional Identity and Input Connectivity of Cortical Afferent Subsets. *Cell Reports*, 31(11):107767.

Kratochwil CF, **Rijli FM** (2020) The Cre/Lox System to Assess the Development of the Mouse Brain. *Methods Mol Biol*, 2047:491-512.

Riva M, Genescu I, Habermacher C, Orduz D, Ledonne F, **Rijli FM**, López-Bendito G, Coppola E, Garel S, Angulo MC, Pierani A. (2019) Activity-dependent death of transient Cajal-Retzius neurons is required for functional cortical wiring. *Elife* 8:e50503.

Coughlan E, Garside VC, Wong SFL, Liang H, Kraus D, Karmakar K, Maheshwari U, **Rijli FM**, Bourne J, McGlinn E. (2019) A Hox Code Defines Spinocerebellar Neuron Subtype Regionalization. *Cell Reports*, 29(8):2408-2421.

Kitazawa T, **Rijli FM** (2018) Barrelette Map Formation in the Prenatal Mouse Brainstem. *Curr Op Neurobiol*, Oct 17;53:210-219.

Kitazawa T, **Rijli FM** (2018) Integrating into the Rhombomere Community across the Border. *Developmental Cell*, 45(5):546-548.

Minoux M, Holwerda S, Vitobello A, Kitazawa T, Kohler H, Stadler M, **Rijli FM** (2017) Gene Bivalency at Polycomb Domains Regulates Cranial Neural Crest Positional Identity. *Science*, 355(6332), eaal2913. (full article)

Kitazawa T, **Rijli FM** (2017) Nuclear Pore Protein Meets Transcription Factor in Neural Fate. *Neuron*, 96(2):259-261.

Karmakar K, Narita Y, Fadok J, Loche A, Fadok J, Genoud C, Ducret S, Di Meglio T, Thierry R, Bacelo J, Lüthi A, **Rijli FM** (2017) Hoxa2 and Hoxb2 Are Required for Tonotopic Map Precision and Sound Discrimination in the Mouse Auditory Brainstem.

Cell Reports, 18(1):185-197.

Kratochwil CF, Maheshwari U, **Rijli FM** (2017) The Long Journey of Pontine Nuclei Neurons: From Rhombic Lip to Cortico-Ponto-Cerebellar Circuitry. *Front Neural Circuits*, 11:33.

Renier N, Dominici C, Erzurumlu RS, Kratochwil CF, **Rijli FM**, Gaspar P, Chédotal A (2017) A mutant with bilateral whisker to barrel inputs unveils somatosensory mapping rules in the cerebral cortex. *Elife*, 6, e23494.

Moreno-Juan V, Filipchuk A, Antón-Bolaños N, Mezzena C, Gezelius H, Andrés B, Rodriguez Malmierca L, Susín R, Schaad O, Rutlin M, Nelson S, Ducret S, Valdeolmillos M, **Rijli FM**, López-Bendito G (2017) Prenatal thalamic waves regulate cortical area size prior to sensory processing. *Nature Communications*, 8:14172.

Daum JM, Keles Ö, Holwerda SJ, Kohler H, **Rijli FM**, Stadler M, Roska B. (2017) The formation of the light-sensing compartment of cone photoreceptors coincides with a transcriptional switch. *Elife*, 6. pii: e31437.

Furutera T, Takechi M, Kitazawa T, Takei J, Yamada T, Vu Hoang T, **Rijli FM**, Kurihara H, Kuratani S, Iseki S. (2017) Differing contributions of the first and second pharyngeal arches to tympanic membrane formation in the mouse and chick. *Development*, 144(18):3315-3324.

Kim M, Minoux M, Piaia A, Kueng B, Gapp B, Weber D, Haller C, Barbieri S, Namoto K, Lorenz T, Wirsching J, Bassilana F, Dietrich W, **Rijli FM**, Ksiazek I. (2017) DPP9 enzyme activity controls survival of mouse migratory tongue muscle progenitors and its absence leads to neonatal lethality due to suckling defect.

Dev Biol, 431(2):297-308.

Laumonnerie C, Bechara A, Vilain N, Kurihara Y, Kurihara H, **Rijli FM** (2015) Facial whisker pattern is not sufficient to instruct a whisker-related topographic map in the mouse somatosensory brainstem. *Development*, 142:3704-3712.

Sreenivasan V, Karmakar K, **Rijli FM***, and Petersen CCH* (2015) Parallel pathways from motor and somatosensory cortex for controlling whisker movements in mice. *Eur J Neurosci* 41, 354-67. *corresponding authors

Lizen B, Claus M, Jeannotte L, **Rijli FM**, Gofflot F (2015) Perinatal induction of Cre recombination with tamoxifen. *Transgenic Res* 24(6), 1065-77.

Kitazawa T, Fujisawa K, Narboux-Nême N, Arima Y, Kawamura Y, Inoue T, Wada Y, Kohro T, Aburatani H, Kodama T, Kim KS, Sato T, Uchijima Y, Maeda K, Miyagawa-Tomita S, Minoux M, **Rijli FM**, Levi G, Kurihara Y, Kurihara H. (2015) Distinct effects of Hoxa2

overexpression in cranial neural crest populations reveal that the mammalian hyomandibular-ceratohyal boundary maps within the styloid process. *Dev Biol*, 402(2):162-74.

Fantini S, Salsi V, Vitobello A, **Rijli FM**, Zappavigna V. (2015) MicroRNA-196b is transcribed from an autonomous promoter and is directly regulated by Cdx2 and by posterior Hox proteins during embryogenesis. *Biochim Biophys Acta*, 1849(8):1066-80.

Kratochwil CF, **Rijli FM** (2014) The Cre/Lox system to assess the development of the mouse brain. *Methods Mol Biol.*, 1082:295-313.

Minoux M, Kratochwil CF, Ducret S, Amin S, Kitazawa T, Kurihara H, Bobola N, Vilain N, **Rijli FM** (2013) Mouse Hoxa2 genetic analysis provides a model for microtia and auricle duplication. *Development*, 140(21):4386-97.

Di Meglio T, Kratochwil CF, Vilain N, Loche A, Vitobello A, Yonehara K, Hrycaj SM, Roska B, Peters AH, Eichmann A, Wellik D, Ducret S, **Rijli FM** (2013) Ezh2 orchestrates topographic tangential migration and connectivity of precerebellar neurons. *Science*, 339(6116):204-7.

Di Bonito M, Narita Y, Avallone B, Sequino L, Mancuso M, Andolfi G, Franzè AM, Puelles L, **Rijli FM***, Studer M* (2013) Assembly of an auditory sensorimotor sub-circuit by a Hox genetic network in the mouse brainstem.

PLoS Genet, 9(2):e1003249. *co-corresponding senior authors

Rhinn M, Miyoshi K, Watanabe A, Kawaguchi M, Ito F, Kuratani S, Baker CV, Murakami Y, **Rijli FM** (2013) Evolutionary divergence of trigeminal nerve somatotopy in amniotes. *J Comp Neurol*, 521(6):1378-94.

Neuronal Migration in the Vertebrate Hindbrain. In: RUBENSTEIN J. L. R. and RAKIC P. (ed.) *Comprehensive Developmental Neuroscience: Cellular Migration and Formation of Neuronal Connections*, volume 2, pp. 377-404 Amsterdam: Elsevier. (book chapter)

Compagnucci C, Debiais-Thibaud M, Coolen M, Fish J, Griffin JN, Bertocchini F, Minoux M, **Rijli FM**, Borday-Birraux V, Casane D, Mazan S, Depew MJ. (2013) Pattern and polarity in the development and evolution of the gnathostome jaw: both conservation and heterotopy in the branchial arches of the shark, *Scyliorhinus canicula*. *Dev Biol*, 377(2):428-48.

Miguez A, Ducret S, Di Meglio T, Parras C, Hmidan H, Haton C, Sekizar S, Mannioui A, Vidal M, Kerever A, Nyabi O, Haigh J, Zalc B, **Rijli FM***, Thomas JL* (2012) Opposing roles for Hoxa2 and Hoxb2 in hindbrain oligodendrocyte patterning. *J Neurosci*, 32(48):17172-85.

*co-corresponding senior authors

Arima Y, Miyagawa-Tomita S, Minoux M, Maeda K, Nishiyama K, Rieko Asai, Ki-Sung Kim KS, Uchijima Y, Ogawa H, **Rijli FM**, Kurihara Y, Kurihara H (2012) Preotic neural crest cells contribute to coronary artery smooth muscle involving endothelin signaling.

Nature Communications, 3:1267.

Pouchelon G, Frangeul L, **Rijli FM**, Jabaudon D. (2012) Patterning of pre-thalamic somatosensory pathways. *Eur J Neurosci*. 35(10):1533-9.

Grainger S, Lam J, Savory JG, Mears AJ, **Rijli FM**, Lohnes D. (2012) Cdx regulates Dll1 in multiple lineages. *Dev Biol.* 361(1):1-11.

Vitobello A, Ferretti E, Lampe X, Vilain N, Ducret S, Ori M, Spetz JF, Selleri L, **Rijli FM** (2011) Hox and Pbx factors control retinoic acid synthesis during hindbrain segmentation. *Developmental Cell*, 20(4):469-82.

Savory JGA, Mansfield M, **Rijli FM** and Lohnes D (2011) Cdx function regulates planar cell polarity. *Development*, 138(7):1361-70.

Beckmann J, Vitobello A, Ferralli J, Brož DK, **Rijli FM**, Chiquet-Ehrismann R. (2011) Human teneurin-1 is a direct target of the homeobox transcription factor EMX2 at a novel alternate promoter. *BMC Dev Biol.* 11:35.

Coppola E, d'Autréaux F, **Rijli FM**, and Brunet JF (2010) Ongoing roles of Phox2 genes during neuronal differentiation. *Development*, 137(24):4211-20.

Erzurumlu, R., Murakami, Y., and **Rijli F.M.** (2010) Mapping the face in the somatosensory brainstem. *Nature Reviews Neuroscience*, 11(4):252-63.

Zhu, P., Narita, Y., Bundschuh, S., Chattopadhyay, B., Arn Bouloires, E., Stepien, A., Arber, S., Sprengel, R., **Rijli F.M.**, and Friedrich R.W (2009) Genetic tools for strong, cell type-selective and conditional gene expression in the zebrafish brain.

Front Neural Circuits, 11;3:21.

Savory JG, Bouchard N, Pierre V, **Rijli FM**, De Repentigny Y, Kothary R, Lohnes D. (2009) Cdx2 regulation of posterior development through non-Hox targets. *Development*, 136(24):4099-110.

Chedotal, A, and **Rijli, F.M.** (2009) Transcriptional control of tangential neuronal migration in the developing forebrain. *Curr Opinion Neurobiol*, 19(2):139-45. (review article)

Narita, Y, and **Rijli FM** (2009) Hox genes in neural patterning and circuit formation in the vertebrate hindbrain. *Curr Topics Dev Biol*, 88, 139-167.

Minoux, M., Antonarakis, G., Kmita, M., Duboule, D., and **Rijli FM** (2009) Rostral and caudal pharyngeal arches share a common neural crest ground pattern. *Development*, 136(4):637-45.

Geisen MJ, Di Meglio T, Pasqualetti M, Ducret S, Brunet JF, Chedotal A, **Rijli FM** (2008) Hox Parologue Group 2 Genes Control the Migration of Mouse Pontine Neurons through Slit-Robo Signaling. *PloS Biology*, 6(6):e142.

Lampe X, Samad OA, Guiguen A, Matis C, Remacle S, Picard JJ, **Rijli FM**, Rezsohazy R. (2008) An ultraconserved Hox-Pbx responsive element resides in the coding sequence of Hoxa2 and is active in rhombomere 4. *Nucleic Acids Res.* 6(10):3214-25.

Chatonnet, F., Wrobel, L.J., Mézières, V., Pasqualetti, M., Ducret, S., Taillebourg, E., Charnay, P., **Rijli, F.M.**, and Champagnat, J. (2007) Distinct roles of Hoxa2 and Krox20 in the development of rhythmic neural networks controlling inspiratory depth, respiratory frequency, and jaw opening. *Neural Development*, 2(1):19-25.

Pasqualetti M., Diaz C., Jean-Sébastien Renaud J-S., **Rijli F.M.***, Glover JC* (2007) Fate-mapping the mammalian hindbrain: Segmental origins of vestibular projection neurons assessed using rhombomere-specific Hoxa2 enhancer elements in the mouse embryo. *J Neurosci*, 27(36):9670-81 *co-corresponding senior authors

Matis C., Oury F., Remacle S., Lampe X., Gofflot F., Picard J.J., **Rijli F.M.***, and Rezsohazy R.* (2007) Identification of Lmo1 as part of a Hox dependent regulatory network for hindbrain patterning. *Dev Dynamics*, 236(9):2675-2684

Takio Y, Kuraku S, Murakami Y, Pasqualetti M, **Rijli FM**, Narita Y, Kuratani S, Kusakabe R. (2007) Hox gene expression patterns in *Lethenteron japonicum* embryos- Insights into the evolution of the vertebrate Hox code. *Dev Biol.*, 308(2):606-20.

Oury, F, and **Rijli F.M.** (2007) Hoxa2: a key gene for the facial somatosensory map. *Med Sci (Paris)*, 23(3):247-9.

Oury, F., Murakami, Y., Renaud, J-S, Pasqualetti, M., Charnay, P., Ren, S-Y., and **Rijli F.M.** (2006) Hoxa2 and rhombomere dependent development of the mouse facial somatosensory map. *Science*, 313(5792):1408-13 (full article)

Sprecher, S.G., Urbach, R., Technau, G.M., **Rijli, F.M.**, Reichert, H. and Hirth F. (2006) The columnar gene vnd is required for tritocerebral neuromere formation during embryonic brain development of *Drosophila*. *Development*, 133(21):4331-9.

Glover, J., Renaud JS, Lampe, X., and **Rijli, F.M.** (2006) Nuclear receptors in hindbrain development in *Advances in Developmental Biology* Elsevier Press, Inc. (book chapter)

Glover JC, Renaud JS, **Rijli FM**. (2006) Retinoic acid and hindbrain patterning. *J Neurobiol.* 66(7):705-25.

Santagati F, Minoux M, Ren SY, and **Rijli F.M.** (2005) Temporal requirement of Hoxa2 in cranial neural crest skeletal morphogenesis. *Development*, 132(22):4927-36.

Baltzinger, M., Ori, M., Pasqualetti, M., Nardi, I., and **Rijli, F.M.** (2005) Hoxa2 knockdown in *Xenopus* results in hyoid to mandibular homeosis. *Dev Dynamics*, 234(4):858-67.

Murakami Y, Uchida K, **Rijli FM**, Kuratani S. (2005) Evolution of the brain developmental plan: Insights from agnathans. *Developmental Biology*, 280(2):249-59.

Abdel Samad O, Geisen MJ, Caronia G, Varlet I, Zappavigna V, Ericson J, Goridis C, and **Rijli FM** (2004) Integration of anteroposterior and dorsoventral regulation of Phox2b transcription in cranial motor neuron progenitors by homeodomain proteins. *Development*, 131:4071-83.

Takio Y, Pasqualetti M, Kuraku S, Hirano S, **Rijli FM** & Kuratani S (2004) Lamprey Hox genes and evolution of jaws. *Nature*, 429, 6989: 1 p following 262

Murakami Y, Pasqualetti M, Takio Y, Hirano S, **Rijli FM**, Kuratani S. (2004) Segmental development of reticulospinal and branchiomotor neurons in lamprey: insights into the evolution of the vertebrate hindbrain. *Development*, 131, 983-95.

Santagati, F., and **Rijli F.M.** (2003). Cranial neural crest and the building of the vertebrate head. *Nature Reviews Neuroscience*, 4, 806–818.

Pattyn, A., Vallstedt, A., Dias, J., Abdel Samad, O., Krumlauf, R., **Rijli, F.M.**, Brunet, J–F, and Ericson, J. (2003) Coordinated Temporal and Spatial Control of Motor Neuron and Serotonergic Neuron Generation from a Common Pool of CNS Progenitors. *Genes & Development*, 17(6):729-37.

Chatonnet F., Domínguez del Toro, E., Thoby-Brisson, M., Champagnat, J., Fortin, G., **Rijli, F.M.**, and Thaëron, C. (2003) From hindbrain segmentation to breathing after birth: developmental patterning in rhombomeres 3 and 4. *Mol Neurobiol.*, 28(3):277-94.

Abdel Samad, O., and **Rijli F.M.** (2003). Genetic Regulation of Hindbrain Development. K.K. Sinha, P. Chandra, eds. *Advances in Clinical Neurosciences* , East Zone Neuro CME, Ranchi, 13, 411–429. (Book chapter)

Pasqualetti, M. and **Rijli, F.M.** (2002). The plastic face. *Nature*, 416, 493–494.

Pasqualetti M, Ren SY, Poulet M, LeMeur M, Dierich A, and **Rijli FM.** (2002) A Hoxa2 knockin allele that expresses EGFP upon conditional Cre-mediated recombination. *Genesis*, 32(2):109-11.

Ren, SY, Angrand, PO, and **Rijli, F.M.** (2002) A targeted insertion results in a rhombomere 2-specific Hoxa2 knockdown and ectopic activation of Hoxa1 expression. *Dev Dynamics*, 225(3):305-15.

Ren SY, Pasqualetti M, Dierich A, Le Meur M, and **Rijli FM.** (2002) A Hoxa2 mutant conditional allele generated by Flp- and Cre-mediated recombination. *Genesis*, 32(2):105-8.

Pasqualetti M., Neun, R., Davenne M., and **Rijli F.M.** (2001) Retinoic acid rescues inner ear defects in Hoxa1 deficient newborn mice. *Nature Genetics*, 29(1):34-9. (full article)

Pasqualetti, M. and **Rijli, F.M.** (2001). Homeobox genes and brainstem developmental disorders: learning from knockout mice. *Curr. Opinion Neurol.*, 14, 177-184.

Dominguez del Toro, E., Borday, V., Davenne, M. Neun, R., **Rijli, F.M.** and Champagnat, J. (2001). Generation of a novel functional neuronal circuit in Hoxa1 mutant mice. *J Neuroscience*, 21(15):5637-42.

Pasqualetti, M. and **Rijli, F.M.** (2001) Vitamin A prevents inner ear defects in mice with congenital homeobox gene deficiency. *TheScientificWorld*, 1, 916–918.

Pasqualetti, M., Ori, M., Nardi, I. and **Rijli, F.M.** (2000). Ectopic Hoxa2 induction after neural crest migration results in homeosis of jaw elements in Xenopus.

Development, 127 (24), 5367-5378.

Davenne, M., Maconochie, M.K., Neun, R., Pattyn, A., Chambon, P., Krumlauf, R., and **Rijli, F.M.** (1999) Hoxa2 and Hoxb2 control dorsoventral patterns of neuronal development in the rostral hindbrain. *Neuron*, 22, 677-691.

Helmbacher, F., Pujades, C., **Rijli, F.M.**, Chambon, P. and Charnay, P. (1998). Hoxa1 and Krox-20 synergize to control the development of rhombomere 3.

Development, 125, 4739-4748.

Gavalas, A., Studer, M., Lumsden, A., **Rijli, F.M.***, Krumlauf, R. and Chambon, P.* (1998). Hoxa1 and Hoxb1 synergize in patterning the hindbrain, cranial nerves and second pharyngeal arch. *Development*, 125, 1123-1136. *co-corresponding senior authors

Studer, M., Gavalas, A., Marshall, H., Ariza-McNaughton, L., **Rijli, F.M.**, Chambon, P., and Krumlauf, R. (1998). Genetic interactions between Hoxa1 and Hoxb1 reveal new roles in regulation of early hindbrain patterning. *Development*, 125, 1025-1036.

Rijli, F.M.*, Gavalas, A., and Chambon, P. (1998). Segmentation and specification in the branchial region of the head: the role of the Hox selector genes. *Int. J. Dev. Biol.*, 42, 393-401.
* corresponding author

Mark, M., **Rijli, F.M.**, and Chambon, P. (1997). Homeobox genes in embryogenesis and pathogenesis. *Pediatric Research*, 42, 421-429.

Rijli, F.M. and Chambon, P. (1997). Genetic interactions of Hox genes in limb development: learning from compound mutants. *Curr. Opinion Genet. Dev.*, 7, 481-487.

Dupé, V., Davenne, M., Brocard, J., Dollé, P., Mark, M., Dierich, A., Chambon, P., and **Rijli, F.M.** (1997). In vivo functional analysis of the Hoxa-1 3' retinoic acid response element (3'RARE). *Development*, 124, 399-410.

Gavalas, A., Davenne, M., Lumsden, A., Chambon, P., and **Rijli, F.M.** (1997). Role of Hoxa-2 in axon pathfinding and rostral hindbrain patterning. *Development*, 124, 3683-3691.

Favier, B., **Rijli, F.M.**, Fromental-Ramain, C., Fraulob, V., Chambon, P., and Dollé P. (1996). Functional cooperation between the non-paralogous genes Hoxa-10 and Hoxd-11 in the developing forelimb and axial skeleton. *Development*, 122, 449-460.

Taneja, R., Thisse, B., **Rijli, F.M.**, Thisse, C., Bouillet, P., Dollé, P., and Chambon, P. (1996). The expression pattern of the mouse receptor tyrosine kinase MDK1 is conserved through evolution and requires Hoxa-2 for rhombomere-specific expression in mouse embryos. *Dev Biol*, 177, 397-412.

Rijli, F.M., Matyas, R., Pellegrini, M., Dierich, A., Gruss, P., Dollé, P., and Chambon, P. (1995). Cryptorchidism and homeotic transformations of spinal nerves and vertebrae in Hoxa-10 mutant mice. *Proc. Natl. Acad. Sci. USA*, 92, 8185-8189.

Mark, M., **Rijli, F.M.**, Lufkin, T., Dollé, P., Gorry, P., and Chambon, P. (1995). Hox gene function and the development of the head. Juurlinck B, Kulik W, Krone P, Verge V, Doucette R, eds, *Neural Cell Specification: Molecular Mechanisms and Neurotherapeutic Implications*, Plenum Press, New York, 3-16. (Book chapter)

Mark, M., **Rijli, F.M.**, and Chambon P. (1995). Alteration of Hox gene expression in the branchial region of the head causes homeotic transformations, hindbrain segmentation defects and atavistic changes. *Seminars in Developmental Biology*, 6, 275-284.

Mark, M., Lohnes, D., Mendelsohn, C., Dupé, V., Vonesch, JL., Kastner, P., **Rijli, F.M.**, Bloch-Zupan, A., Chambon, P. (1995). Roles of retinoic acid receptors and of Hox genes in the patterning of the teeth and of the jaw skeleton. *Int. J. Dev. Biol.*, 39, 111-121.

Rijli, F.M., Dollé, P., Fraulob, V., LeMeur, M., and Chambon, P. (1994). Insertion of a targeting construct in a Hoxd-10 allele can influence the control of Hoxd-9 expression. *Dev Dynamics*, 201, 366-377.

Rijli, F.M., Mark, M., Lakkaraju, S., Dierich, A., Dollé, P., and Chambon, P. (1993). A homeotic transformation is generated in the rostral branchial region of the head by disruption of Hoxa-2, which acts as a selector gene. *Cell*, 75, 1333-1349.

Rijli, F.M., De Lucchini, S., Ciliberto, G. and Barsacchi, G. (1993). A Zn-finger protein, Xfin, is expressed during cone differentiation in the retina of the frog *Xenopus laevis*. *Int. J. Dev. Biol.*, 37, 311-317.

De Lucchini, S., **Rijli, F.M.**, Ciliberto, G. and Barsacchi, G. (1991). A Xenopus multifinger protein, Xfin, is expressed in specialized cell types and is localized in the cytoplasm. *Mech. Dev.* 36, 31-40.

Vignal, R., **Rijli, F.M.**, Batistoni, R., Fratta, D., Cremisi, F. and Barsacchi, G. (1991) Two dispersed highly repeated DNA families of *Triturus vulgaris meridionalis* (Amphibia, Urodela) are widely conserved among Salamandridae. *Chromosoma*, 100, 87-96.

Moore, S.K., **Rijli, F.** and Appella, E. (1990). Characterization of the mouse 84 kDa heat shock protein gene family. *DNA and Cell Biology* 9, 387-400.